Crossroads of America Council BSA Annual BSA Health and Medical Record
Part A GENERAL INFORMATION
Name
Address

Crossroads of America Council BSA Annual BSA Health and Medical Record Part A GENERAL INFORMATION			High-adventure base participants: Expedition/crew No.: or staff position:				
Name				Age		☐ Female ☐	
				Grade	e completed (youth only)		
Unit leade	r		Council name/No.		Unit No		
Social S	Security 1	No. (optional; may be required by medical facilities for	treatment)	Religious preference	ce		
Health/acc	ident insur	ance company	Policy No.				
n case Name		gency, notify:	Relatio	onship			
Address							
Home ph	one	Business	phone	Cell phone			
Alternate c	contact		Alternate's	s phone			
HEALTH	HISTORY	′					
Are you	now, or h	nave you ever been treated for any of the following	:	Aller	gies or Reaction	to:	
Yes	No	Condition	Explain	Medication			
		Asthma Last attack:		Food, Plants, or	Insect Bites		
		Diabetes Last HbA1c:					
		Hypertension (high blood pressure)					

Yes	No	Condition	E	cplain	Medica	tion	
		Asthma Last attack:			Food,	Plants,	or Insect Bites
- 1		Diabetes Last HbA1c:			7		
		Hypertension (high blood pressure)			7		Immunizations:
- 44		Heart disease (e.g., CHF, CAD, MI)			The f	ollowin	g are recommended by the BSA.
- 8		Stroke/TIA					munization is required and mu
		Lung/respiratory disease					eceived within the last 10 years. If ha
		Ear/sinus problems					It "D" and the year. If immunize ox and the year received.
		Muscular/skeletal condition			1		Date
		Menstrual problems (women only)			3200		
		Psychiatric/psychological and emotional difficulties					Tetanus
		Behavioral disorders (e.g., ADD, ADHD, Asperger syndrome, autism)					Diphtheria
		Bleeding disorders					Mumps
3		Fainting spells					Rubella
		Thyroid disease					Polio
- 8		Kidney disease	4				Chicken pox
		Sickle cell disease					Hepatitis A
- 6		Seizures Last seizure:	3				Hepatitis B
- 8		Sleep disorders (e.g., sleep apnea)	Use CPAP: Yes □	No 🗆	4.774		Influenza
		Abdominal/digestive problems		5 5592	1 -	100000	
- 3		Surgery					Other (i.e., HIB)
		Serious injury] 🗆 Ex	emptic	on to immunizations claimed
- 3		Other			(fo	rm req	uired).

MEDICATIONS

List all medications currently used. (If additional space is needed, please photocopy this part of the health form.) Inhalers and EpiPen information must be included, even if they are for occasional or emergency use only.

(For more information about immunizations, as well as the immunization exemption form, see Scouting Safely on Scouting.org.)

Medication Frequency Approximate date started Reason for medication	Medication Frequency Approximate date started Reason for medication	Medication Frequency Approximate date started Reason for medication	
Medication Strength Frequency Approximate date started Reason for medication	Medication Frequency Approximate date started Reason for medication	Medication Frequency Approximate date started Reason for medication	

Administration of the above medications is approved by (if required by your state): _

Parent/guardian signature and/or MD/DO, NP, or PA signature

Be sure to bring medications in sufficient quantities and the original containers. Make sure that they are NOT expired, including inhalers and EpiPens. You SHOULD NOT STOP taking any maintenance medication.

Part B

Part B

Full name:

INFORMED CONSENT AND HOLD HARMLESS/RELEASE AGREEMENT

High-adventure base participants:
Expedition/crew No.:
or staff position:

I understand that participation in Scouting activities involves a certain degree of risk and can be physically, mentally, and emotionally demanding. I also understand that participation in these activities is entirely voluntary and requires participants to abide by applicable rules and standards of conduct.

In case of an emergency involving me or my child, I understand that every effort will be made to contact the individual listed as the emergency contact person. In the event that this person cannot be reached, permission is hereby given to the medical provider selected by the adult leader in charge to secure proper treatment, including hospitalization, anesthesia, surgery, or injections of medication for me or my child. Medical providers are authorized to disclose protected health information to the adult in charge, camp medical staff, camp management, and/or any physician or health care provider involved in providing medical care to the participant. Protected Health Information/Confidential Health Information (PHI/CHI) under the Standards for Privacy of Individually Identifiable Health Information, 45 C.F.R. §§160.103, 164.501, etc. seq., as amended from time to time, includes examination findings, test results, and treatment provided for purposes of medical evaluation of the participant, follow-up and communication with the participant's parents or guardian, and/or determination of the participant's ability to continue in the program activities.

I have carefully considered the risk involved and give consent for myself and/or my child to participate in these activities. I approve the sharing of the information on this form with BSA volunteers and professionals who need to know of medical situations that might require special consideration for the safe conducting of Scouting activities.

require special continuoration for the care contacting of cocaling detrines.		
I release the Boy Scouts of America, the local council, the activity coordinator organizations associated with the activity from any and all claims or liability arising		er
☐ Without restrictions.		
☐ With special considerations or restrictions (list)		
TALENT RELEASE AGREEMENT		
I hereby assign and grant to the local council and the Boy Scouts of America the rigilm/videotapes/electronic representations and/or sound recordings made of me release the Boy Scouts of America, the local council, the activity coordinators, ar organizations associated with the activity from any and all liability from such use	or my child at all Scouting activities, and I hereby all employees, volunteers, related parties, or other	
I hereby authorize the reproduction, sale, copyright, exhibit, broadcast, elect film/videotapes/electronic representations and/or sound recordings without lim and I specifically waive any right to any compensation I may have for any of the	nitation at the discretion of the Boy Scouts of America,	
ADULTO ALITHODIZED TO TAKE VOLITILITO AND EDOM EVENTO.		
ADULTS AUTHORIZED TO TAKE YOUTH TO AND FROM EVENTS: You must designate at least one adult. Please include a telephone number.		
Name 1. Name	Tolophono	
2. Name	•	
3. Name	•	dults
NOT authorized to take youth to and from events:		adito
1. Name		
2. Name		
3. Name		
I understand that, if any information I/we have provided is found to be inac participation in any event or activity. If I am participating at Philmont, Philmont Training Center, Northern Ti understand the risk advisories explained in Part D, including height and that the participant will not be allowed to participate in applicable high-a The participant has permission to engage in all high-adventure activities health-care provider.	ccurate, it may limit and/or eliminate the opportunity fier, or Florida Sea Base: I have also read and weight requirements and restrictions, and understadventure programs if those requirements are not m	nd et.
Participant's name		
Participant's signature		
Parent/guardian's signature(if participant is under	Date	
This Annual Health and Medical Record is valid for 12 calendar months.	uie age oi 10)	

680-001 2010 Printing Rev. 11/2010

DOB:

	High-adventu Expedition/crew or staff position:	•	articipants:		
TO THE EXAMINING HEALTH-CARE PROVIDER (Certification re being asked to certify that this individual has no contraindication diventure program at one of the national high-adventure bases, please Part D was made available to me. Yes No)	for participation	in a Scoutir	ng experienc	e. For individuals	
HYSICAL EXAMINATION					
leight (inches) Weight (pounds) M	laximum weight	for height _	N	leets height/weigh	nt lindats Yes No
lood pressurePulse					
If you exceed the maximum weight for height as explained on this p away from an emergency vehicle-accessible roadway, you will not and/or camp, participation of an individual exceeding the maximum health-care provider is determined to be 20 percent or less for a femtest to be used for this determination.) Please call the event leader a strongly encouraged for all other events.	be allowed to pa weight for heigh ale or 15 percent	articipate. At t may be all t or less for	t the discretion lowed if the b a male. (Phili	on of the medical body fat percentag mont requires a w	advisors of the event ge measured by the vater-displacement
Normal Abnormal Explain Any Abnormalities	Range of	Mobility	Normal	Abnormal	Explain Any Abnormalities
Eyes	Knees (both)				
Ears	Ankles (both)				
Nose	Spine				
Throat					
Lungs					
Neurological	Othe	r	Yes	No	
Heart	Contacts				
Abdomen	Dentures				
Genitalia	Braces				
Skin	Inguinal hernia				Explain
Emotional	Medical equipm (i.e., CPAP, ox)	nent			
adjustment	□ Neg		Positive	•	
Ilergies (to what agent, type of reaction, treatment):					
estrictions (if none, so state)					
XAMINER'S CERTIFICATION	Height (inches)	Recomm Weight		Allowable Exception	Maximum Acceptance
certify that I have reviewed the health history and examined this person nd find no contraindications for participation in a Scouting experience.	60	97-	` '	139-166	166
his participant	61	101-	-143	144-172	172
Meets height/weight requirements Does not have uncontrolled heart disease, asthma, or hypertension	62	104-		149-178	178
Has not had an orthopedic injury, musculoskeletal problems, or	63 64	107-		153-183 158-189	183 189
orthopedic surgery in the last six months or possesses a letter of	65	114-		163-195	195
clearance from their orthopedic surgeon or treating physician • as no uncontrolled psychiatric disorders	66	118-		168-201	201
Has had no seizures in the last year	67	121-		173-207	207
Does not have poorly controlled diabetes	68	125- 129-		179-214 186-220	214
If less than 18 years of age and planning to scuba dive, does not have diabetes, asthma, or seizures	70	132-		189-226	226
ovider printed name	71	136-		195-233	233
OTION PIRROUTIGITIO	72	140-		200-239	239
dress	73	144-		206-246	246
y, state, zip	74 75	148-21 152-21		211-252 217-260	252 260
<i>y</i> , σαιο, ωρ	75 76	152-21		217-260	267
ice phone	77	160-22		229-274	274
gnature	78	164-		235-281	281
, MINIO	79 & over	170-	-240	241-295	295
te	This table is based Dept. of Agriculture	on the revised Dand the Dent of	Dietary Guidelines Health & Human	for Americans from the Services.	U.S.
DO NOT V	VRITE IN THIS B		James G. Milliott		
REVIEW FOR CAMP OR SPECIAL ACTIVITY	VICITE III II III D	UX .			
	WILL IN THIS D		Da	ate	

Part C Full name: ______DOB: _____

Part D

Participation at any of the BSA's high-adventure bases can be physically, mentally, and emotionally demanding. To be better prepared, each participant must complete the following before attending any high-adventure base:

- Fill in parts A and B of the Annual Health and Medical Record.
- Share Part D with the examining health-care provider.
- Have a physical exam by a certified and licensed health care provider/physician (MD, DO), nurse practitioner, or physician assistant, and have part C completed.
- Read the following information, which focuses on specific risks at the high-adventure base you will be attending.

The Trek Experience. Each high-adventure base offers a unique experience that is not risk-free. Knowledgeable staff will instruct all participants in safety measures to be followed. Be prepared to listen to and carefully follow these safety measures and to accept responsibility for the health and safety of yourself and others.

Philmont. Each participant must be able to carry a 35- to 50-pound pack while hiking 5 to 12 miles per day in an isolated mountain wilderness ranging from 6,500 to 12,500 feet in elevation. Summer/autumn climatic conditions include temperatures from 30 to 90 degrees, low humidity (10 to 30 percent), and frequent, sometimes severe, afternoon thunderstorms. Activities include horseback riding, rock climbing and rappelling, challenge events, pole climbing, black powder shooting, 12-gauge trap shooting, .30-06 shooting, trail building, mountain biking, and other activities that have potential for injury.

Winter climatic conditions can range from -20 to 60 degrees. For the Kanik Experience, each person will walk, ski, or snowshoe along snow-covered trails pulling loaded toboggans or sleds for up to 3 miles, or more on a cross-country ski trek. Refer to the Philmont Scout Ranch website for specific information.

Northern Tier. Each person must be able to carry a 50- to 85-pound pack or canoe from a quarter-mile to 2 miles several times a day on rough, swampy, and rocky portages and paddle 10 to 15 miles per day, often against a headwind. Climatic conditions can range from 30 to 100 degrees in summer/autumn and from -40 to 40 degrees in the winter. For the Okpik Experience, each person will walk, ski, or snowshoe along snow-covered trails or across frozen lakes, pulling loaded toboggans or sleds for up to 3 miles, or more if on a cross-country ski trek. Refer to the Northern Tier website for specific information.

Florida Sea Base. Climatic conditions at Florida Sea Base include temperatures ranging from 50 to 95 degrees, high humidity, heat index reaching to 110 degrees, and frequent, sometimes severe, afternoon thunderstorms. Activities include snorkeling, scuba diving, kayaking, canoeing, sailing, hiking, and other activities that have potential for injury. Refer to the Sea Base website for specific information.

Risk Advisory. All of the high-adventure bases have excellent health and safety records and strive to minimize risks to participants and advisors by emphasizing appropriate safety precautions. Because most participants are prepared, are conscious of risks, and take safety precautions, they do not experience injuries. If you decide to attend Philmont, Northern Tier, or Florida Sea Base, you should be physically fit, have proper clothing and equipment, and be willing to follow instructions, work as a team with your crew, and take responsibility for your own health and safety.

Parents, guardians, and participants in any high-adventure program are advised that journeying to and from these bases can involve exposure to accidents, illness, and/or injury.

High-adventure staff members have been trained in first aid, CPR, and accident prevention and are prepared to assist the adult advisor in recognizing, reacting to, and responding to accidents, injuries, and illnesses as needed. Each crew is required to have at least one member trained in wilderness first aid and CPR. Medical and search-and-rescue services are provided in response to an accident or emergency. However, response times can be affected by location, terrain, weather, or other emergencies and could be delayed for hours or even days in a wilderness setting.

Philmont. Participants and guests for Philmont activities that are conducted with limited access to the backcountry, including most Philmont Training Center conferences and family programs, should review Part D to understand potential health risks inherent at 6,700 feet in elevation in a dry Southwest environment.

High elevation; physically demanding high-adventure program in remote mountainous areas; camping while being exposed to occasional severe weather conditions such as lightning, hail, flash floods, and heat; and other potential problems, including injuries from tripping and falling, falls from horses, heat exhaustion, and motor vehicle accidents, can worsen underlying medical conditions. Philmont's trails are steep and rocky. Wild animals such as bears, rattlesnakes, and mountain lions are native and usually present little danger if proper precautions are taken. Please call Philmont (575-376-2281) if you have any questions.

Northern Tier. While participating in Northern Tier's canoeing and camping wilderness areas, life jackets must be worn at all times when on the water. Crew members travel together at all times. Emergency communications via radio, and in more remote locations by satellite phone, are provided by Northern Tier. Radio communication and/ or emergency evacuation can be hampered by weather, terrain, distance, equipment malfunction, and other factors, and are not a substitute for taking appropriate precautions and having adequate first-aid knowledge and equipment. Please call Northern Tier (218-365-4811) if you have any questions.

Florida Sea Base. Several activities are offered, including snorkeling, sailing, camping, kayaking, canoeing, swimming, fishing, and scuba diving. Diving is an exciting and demanding activity. When performed correctly, it is very safe. When established safety procedures are not followed, however, there are extreme dangers. All participants will need to learn from the instructor the important safety rules regarding breathing and equalization while scuba diving. Improper use of scuba equipment can result in serious injury, so participants must be instructed to use the equipment safely under direct supervision of a qualified instructor.

To scuba dive safely, participants must not be extremely overweight or in poor physical condition. Diving can be strenuous under certain conditions. Participants' respiratory and circulatory systems must be in good health. All body air spaces must be normal and healthy. A person with heart trouble, a current cold or congestion, epilepsy, asthma, or a severe medical problem, or who is under the influence of alcohol or drugs, should not dive. If taking medication, participants should consult a doctor and the instructor before participation in this program. If there is a question about the advisability of participation, contact the family physician first, then call the Sea Base at 305-664-4173. The Sea Base health supervisor reserves the right to make medical decisions regarding the participation of individual at Sea Base.

Food. Each base offers food appropriate for the experience. If a participant has a problem with the diet described in the participant guide, please contact the high-adventure base you are considering attending.

Medications. Each participant who has a condition requiring medication should bring an appropriate supply for the duration of the trip. Consider bringing duplicate or even triplicate supplies of vital medications. People with allergies that have resulted in severe reactions or anaphylaxis must bring with them an EpiPen that has not expired.

Immunizations. Each participant must have received a tetanus immunization within the last 10 years. Recognition will be given to the rights of those Scouts and Scouters who do not have immunizations because of philosophical, political, or religious beliefs. In such a situation, the Immunization Exemption Request form is required.

Recommendations Regarding Chronic Illnesses. Each base requires that this information be shared with the parents or guardians and examining physician of every participant. There are no facilities for extended care or treatment; therefore participants who cannot meet these requirements will be sent home at their expense.

Staff and/or staff physicians reserve the right to deny the participation of any individual on the basis of a physical examination and/or medical history.

Adults or youth who have had any of the following conditions should undergo a thorough evaluation by a physician before considering participation at a BSA high-adventure base.

Cardiac or Cardiovascular Disease, including:

- 1. Angina (chest pain caused by blocked blood vessels or coming from the heart)
- 2. Myocardial infarction (heart attack)
- 3. Heart surgery or heart catheterization (including angioplasty to treat blocked blood vessels, balloon dilation, or stents)
- 4. Stroke or transient ischemic attacks (TIAs)
- 5. Claudication (leg pain with exercise, caused by hardening of the arteries)
- 6. Family history of heart disease or a family member who died unexpectedly before age 50
- 7. Diabetes
- 8. Smoking
- 9. Excessive weight

Youth who have a congenital heart disease or an acquired heart disease such as rheumatic fever, Kawasaki's disease, or mitral valve prolapse should undergo thorough evaluation by a physician before considering participating at a high-adenture base. The physical exertion at any of the high-adventure bases may precipitate either a heart attack or stroke in susceptible persons. Participants with a history of any of the first seven conditions listed above should have a physician-supervised stress test. More extensive testing (e.g., nuclear stress test) is recommended for participants who have coronary heart disease. Even if the stress test results are normal, the results of testing done at lower elevations, without backpacks, do not guarantee safety. If the test results are abnormal, the individual is advised not to participate.

Hypertension (High Blood Pressure). The combination of physical, mental, and emotional stress, increased exertion and/or heat, and altitude appears to cause a significant increase in blood pressure in some individuals. Occasionally, hypertension reaches such a level that it is no longer safe to engage in strenuous activity. Hypertension can increase the risk of having a stroke, heart attack, or angina. Participants should have a normal blood pressure (less than 140/90). Persons with significant hypertension (greater than 140/90) should be treated and controlled before attending any highadventure base, and should continue on medications while participating. The goal of treatment should be to lower the blood pressure to normal. Participants already on antihypertensive therapy with normal blood pressure should continue on medications. Individuals taking diuretics to treat hypertension are at increased risk for dehydration related to strenuous physical activity and should be careful to maintain good hydration during the trek.

Philmont. Each participant who is 18 years of age or older will have his or her blood pressure checked at Philmont. Those individuals with a blood pressure consistently greater than 160/100 at Philmont may be kept off the trail until their blood pressure decreases.

Florida Sea Base. Those taking beta-blocker medication should consider a change of medication before participating in any scuba program.

Insulin-Dependent Diabetes Mellitus. Exercise and the type of food eaten affect insulin requirements. Any individual with insulin-dependent diabetes mellitus should be able to monitor personal blood glucose and to know how to adjust insulin doses based on these factors. The person with diabetes also should know how to give a self-injection. Both the person with diabetes and one other person in the group should be able to recognize indications of excessively high blood sugar (hyperglycemia or diabetic ketoacidosis) and excessively low blood sugar (hypoglycemia). The person with diabetes and one other individual should know the appropriate initial responses for these conditions. An insulin-dependent

person who has been newly diagnosed (within the last six months) or who has undergone a change in delivery system (e.g., insulin pump) in the last six months should not attempt to participate. A person with diabetes who has had frequent hospitalizations for diabetic ketoacidosis or who has had frequent problems with hypoglycemia should not participate until better control of the diabetes has been achieved.

Philmont. It is recommended that the person with diabetes and one other individual carry insulin on the trek (in case of accident) and that a third vial be kept at the Health Lodge for backup. Bring insulin in a small insulated container. Bring enough testing equipment and supplies for the entire trip and trek. Extras are usually needed. If an individual has been hospitalized for diabetes-related illnesses within the past year, the individual must obtain permission to participate by contacting the Philmont Health Lodge at 575-376-2281.

Florida Sea Base. Persons with diabetes who are 18 years of age or older who wish to scuba dive should be assessed by a physician familiar with both hyperbaric issues related to diabetes and medications used for the control and treatment of diabetes. Persons 18 years old or older who are determined to be candidates for scuba diving must submit four hemoglobin A1c (HbA1c) tests, each with HbA1c values less than 7, taken within the previous 12 months. Any test within the past 12 months with an HbA1c value greater than 7 disqualifies a person from scuba diving as part of a BSA activity.

Persons less than 18 years of age with Type 1 diabetes will not be allowed to scuba dive. Persons under the age of 18 who control their diabetes with exercise and diet (no medications) and can provide three sequential hemoglobin tests with HbA1c values less than 6 may be approved to scuba dive.

<u>Seizures (Epilepsy)</u>. A seizure disorder or epilepsy does not exclude an individual from participating at a high-adventure base. However, the seizure disorder should be well-controlled by medications. A minimum one year seizure-free period is considered to be adequate control. Exceptions to this guideline may be considered on an individual basis, and will be based on the specific type of seizure and the likely risks to the individual and to other members of the crew.

Florida Sea Base. Any seizure activity within the past five years, regardless of control and/or medication, disqualifies an individual from participation in any scuba program. A person with a history of seizure activity who has been asymptomatic AND medication-free for five years, as evidenced by a physician, will be allowed to dive.

Asthma. Asthma should be well-controlled before participating at any high-adventure base. Well-controlled asthma means: 1) the use of a rescue inhaler (e.g., albuterol) less than once daily; 2) no need for nighttime treatment with a rescue inhaler (e.g., albuterol). Well-controlled asthma may include the use of long-acting bronchodilators, inhaled steroids, or oral medications such as Singulair. You must meet these guidelines in order to participate. You will not be allowed to participate if: 1) you have exercise asthma not prevented by medications; or 2) you have been hospitalized or have gone to the emergency room to treat asthma in the past six months; or 3) you have needed treatment with intravenous, intramuscular, or oral steroids (prednisone) in the past six months. You must bring an ample supply of your medications and a spare rescue inhaler that are not expired. At least one other member of the crew should know how to recognize signs of worsening asthma or an asthma attack, and should know how to use the rescue inhaler. Any person who has needed treatment for asthma in the past three years must carry a rescue inhaler on the trek. If you do not bring a rescue inhaler, you must buy one before you will be allowed to participate.

Florida Sea Base. Persons being treated for asthma (including reactive airway disease) are disqualified from BSA scuba programs. Persons with a history of asthma who have been asymptomatic and have not used medications to control asthma for five years or more may be allowed to scuba dive as part of a BSA activity upon submission of evidence from their treating physician. Persons with a history of asthma who have been asymptomatic and have not used medication to control asthma for less than five years may be allowed to scuba dive as part of a BSA activity upon submission of a methacholine challenge test showing the asthma to be resolved.

Allergy or Anaphylaxis. Persons who have had an anaphylactic reaction from any cause must contact the high-adventure base before arrival. If you are allowed to participate, you will be required to have appropriate treatment with you. You and at least one other member of your crew must know how to give the treatment. If you do not bring appropriate treatment with you, you will be required to buy it before you will be allowed to participate.

Recent Musculoskeletal Injuries and Orthopedic Surgery. Every participant will put a great deal of strain on feet, ankles, and knees due to negotiating steep, rocky trails with a backpack; paddling and portaging heavy gear over irregular terrain; or climbing into and out of a boat. Therefore, individuals with significant musculoskeletal problems (including back problems) or orthopedic surgery/injuries within the last 6 months must have a letter of clearance from their orthopedic surgeon or treating physician to be considered for approval to participate. Permission is not guaranteed. A person with a cast on any extremity may participate only if approved by the high-adventure base. Ingrown toenails are a common problem and must be treated 30 days prior to arrival.

Psychological and Emotional Difficulties. A psychological disorder does not necessarily exclude an individual from participation. Parents and advisers should be aware that no high-adventure experience is designed to assist participants in overcoming psychological or emotional problems. Experience demonstrates that these problems frequently become magnified, not lessened, when a participant is subjected to the physical and mental challenges of a remote wilderness setting. Any condition should be well-controlled without the services of a mental health practitioner. Under no circumstance should medication be stopped immediately prior to participation, and medication should be continued throughout the entire high-adventure experience. Participants requiring medication must bring an appropriate supply for the duration of the trip.

Weight Limits. Weight limit guidelines are used because overweight individuals are at a greater risk for heart disease, high blood pressure, stroke, altitude illness, sleep problems, and injury. Those who fall within the limits are more likely to have an enjoyable trek and avoid incurring health risks.

Philmont. Each participant in a Philmont trek must not exceed the maximum acceptable limit in the weight chart shown below. The right-hand column shows the maximum acceptable weight for a person's height in order to participate in a Philmont trek.

Participants 21 years and older who exceed the maximum acceptable weight limit for their height at the Philmont medical recheck WILL NOT be permitted to backpack or hike at Philmont. They will be sent home. For example, a person 70 inches tall cannot weigh more than 226 pounds. All heights and weights will be measured in stocking feet.

For participants under 21 years of age who exceed the maximum acceptable weight for height, the Philmont physicians will use their best professional judgment in determining participation in a Philmont trek. Philmont will consider up to 20 pounds over the maximum acceptable as stated on the chart; however exceptions are not made automatically, and discussion in advance with Philmont is required regarding any exception to the weight limit for persons under 21 years of age. Philmont's telephone number is 575-376-2281.

Under no circumstances will any individual weighing more than 295 pounds be permitted to participate in backcountry programs. This requirement is necessary due to rescue equipment restrictions and for the safety of search-and-rescue personnel.

The maximum weight for any participant in a Cavalcade Trek and for horse rides is 200 pounds.

Participants and guests in Philmont activities, including most Philmont Training Center conference and family programs, who will participate in limited backcountry access during their visit must not exceed the maximum acceptable limit in the weight chart.

Northern Tier. Each participant in a Northern Tier expedition should not exceed the maximum acceptable weight for height in the table shown on the Annual Health and Medical Record form. Those who fall within the recommended weight limits are much more likely to have an enjoyable trek and avoid incurring injuries and health risks. Extra weight puts strain on the back, joints, and feet. The portage trails can be very muddy, slippery, and rocky, and present a potential for tripping and falling. We also strongly recommend that no participant be less than 100 pounds in weight. Extremely small participants will have a very difficult time carrying canoes and heavy packs.

Canoes' loads are another important reason to limit participant weight. Northern Tier assigns three people to a canoe. The total participant load per canoe must not exceed 600 pounds, or an average of 200 pounds per participant. Northern Tier does not permit individuals exceeding 295 pounds to participate in high-adventure programs.

Florida Sea Base. Any participant or advisor who exceeds the maximum weight limits on the weight chart may want to reconsider participation in a Sea Base high-adventure program. Anyone who exceeds these limits is at extreme risk for health problems. Participants who fall within the guidelines are more likely to have an enjoyable program and avoid incurring health risks. The absolute weight limit for our programs is 295 pounds.

Height (inches)	Recommended Weight (lbs)	Allowable Exception	Maximum Acceptance
60	97-138	139-166	166
61	101-143	144-172	172
62	104-148	149-178	178
63	107-152	153-183	183
64	111-157	158-189	189
65	114-162	163-195	195
66	118-167	168-201	201
67	121-172	173-207	207
68	125-178	179-214	214
69	129-185	186-220	220

Height (inches)	Recommended Weight (lbs)	Allowable Exception	Maximum Acceptance
70	132-188	189-226	226
71	136-194	195-233	233
72	140-199	200-239	239
73	144-205	206-246	246
74	148-210	211-252	252
75	152-216	217-260	260
76	156-222	223-267	267
77	160-228	229-274	274
78	164-234	235-281	281
79 & over	170-240	241-295	295

This table is based on the revised Dietary Guidelines for Americans from the U.S. Dept. of Agriculture and the Dept. of Health & Human Services.









Participant Record (Confidential Information)

Please read carefully before signing.

This is a statement in which you are informed of some potential risks involved in scuba diving and of the conduct required of you during the scuba training program. Your signature on this statement is required for you to participate in the scuba training program offered

by	and
Instructor	
	located in the
Facility	
city of, state/province of _	·
Read this statement prior to signing it. You Medical Statement, which includes the medical quest enroll in the scuba training program. If you are a mithis Statement signed by a parent or guardian. Diving is an exciting and demanding activity.	tionnaire section, to nor, you must have

Diving is an exciting and demanding activity. When performed correctly, applying correct techniques, it is relatively safe. When

Divers Medical Questionnaire To the Participant:

The purpose of this Medical Questionnaire is to find out if you should be examined by your doctor before participating in recreational diver training. Apositive response to a question does not necessarily disqualify you from diving. Apositive response means that there is a preexisting condition that may affect your safety while diving and you must seek the advice of your physician prior to engaging in dive activities

while diving and you must seek the advice of your physician prior to engaging in dive activities.
Could you be pregnant, or are you attempting to become pregnant?
Are you presently taking prescription medications? (with the exception of birth control or anti-malarial)
Are you over 45 years of age and can answer YES to one or more of the following? • currently smoke a pipe, cigars or cigarettes • have a high cholesterol level • have a family history of heart attack or stroke • are currently receiving medical care • high blood pressure • diabetes mellitus, even if controlled by diet alone
Have you ever had or do you currently have
Asthma, or wheezing with breathing, or wheezing with exercise?
Frequent or severe attacks of hayfever or allergy?
Frequent colds, sinusitis or bronchitis?
Any form of lung disease?
Pneumothorax (collapsed lung)?
Other chest disease or chest surgery?
Behavioral health, mental or psychological problems (Panic attack, fear of closed or open spaces)?
Epilepsy, seizures, convulsions or take medications to prevent them?
Recurring complicated migraine headaches or take medications to prevent them?
Blackouts or fainting (full/partial loss of consciousness)?

Frequent or severe suffering from motion sickness (seasick, carsick,

established safety procedures are not followed, however, there are increased risks.

To scuba dive safely, you should not be extremely overweight or out of condition. Diving can be strenuous under certain conditions. Your respiratory and circulatory systems must be in good health. All body air spaces must be normal and healthy. A person with coronary disease, a current cold or congestion, epilepsy, a severe medical problem or who is under the influence of alcohol or drugs should not dive. If you have asthma, heart disease, other chronic medical conditions or you are taking medications on a regular basis, you should consult your doctor and the instructor before participating in this program, and on a regular basis thereafter upon completion. You will also learn from the instructor the important safety rules regarding breathing and equalization while scuba diving. Improper use of scuba equipment can result in serious injury. You must be thoroughly instructed in its use under direct supervision of a qualified instructor to use it safely.

If you have any additional questions regarding this Medical Statement or the Medical Questionnaire section, review them with your instructor before signing.

Please answer the following questions on your past or present medical history
with a YES or NO. If you are not sure, answer YES. If any of these items apply
to you, we must request that you consult with a physician prior to participating in
scuba diving. Your instructor will supply you with an RSTC Medical Statement
and Guidelines for Recreational Scuba Diver's Physical Examination to take to your physician.
Dysentery or dehydration requiring medical intervention?
Any diverse idente en decembra element

Dysentery or dehydration requiring medical intervention?
Any dive accidents or decompression sickness?
Inability to perform moderate exercise (example: walk 1.6 km/one mile within 12 mins.)?
Head injury with loss of consciousness in the past five years?
Recurrent back problems?
Back or spinal surgery?
Diabetes?
Back, arm or leg problems following surgery, injury or fracture?
High blood pressure or take medicine to control blood pressure?
Heart disease?
Heart attack?
Angina, heart surgery or blood vessel surgery?
Sinus surgery?
Ear disease or surgery, hearing loss or problems with balance?
Recurrent ear problems?
Bleeding or other blood disorders?
Hemia?
Ulcers or ulcer surgery ?
Acolostomy or ileostomy?
Recreational drug use or treatment for, or alcoholism in the past five

The information I have provided about my medical history is accurate to the best of my knowledge. I agree to accept responsibility for omissions regarding my failure to disclose any existing or past health condition.

years?

STUDENT

Please print legibly.							
Name	Initial	Last	Bi	rth Date	v/Month/Year	Age	
Mailing Address							
City				ce/Region			
Country			Zip/Postal C	Ode			
Home Phone ()		Business F	hone ()				
Email			FAX				
Name and address of your family p	hysician						
Physician			Clinic/Hospital				
Address							Date
of last physical examination							
Name of examiner			Clinic/Hospital				
Address							
Phone ()		Email					
PHYSICIAN							
This person applying for training capplicant's medical fitness for scul						diving. Your opir	nion of the
Physician's Impression							
☐ I find no medical conditions	s that I consider incom	patible with diving.	I				
☐ am unable to recommend the	his individual for diving.						
Remarks							
				Date			
Physician's Signature or Legal Re	epresentative of Medical Practitione	er		<i></i>	Da	y/Month/Year	
Physician			Clinic/Hospital				
Address							
Phone ()		Email					

Guidelines for Recreational Scuba Diver's Physical Examination

Instructions to the Physician:

Recreational **SCUBA** (Self-Contained Underwater Breathing Apparatus) can provide recreational divers with an enjoyable sport safer than many other activities. The risk of diving is increased by certain physical conditions, which the relationship to diving may not be readily obvious. Thus, it is important to screen divers for such conditions.

The RECREATIONAL SCUBA DIVER'S PHYSICAL EXAMINA-

TION focuses on conditions that may put a diver at increased risk for decompression sickness, pulmonary overinflation syndrome with subsequent arterial gas embolization and other conditions such as loss of consciousness, which could lead to drowning. Additionally, the diver must be able to withstand some degree of cold stress, the physiological effects of immersion and the optical effects of water and have sufficient physical and mental reserves to deal with possible emergencies.

The history, review of systems and physical examination should include as a minimum the points listed below. The list of conditions that might adversely affect the diver is not all-inclusive, but contains the most commonly encountered medical problems. The brief introductions should serve as an alert to the nature of the risk posed by each medical problem.

The potential diver and his or her physician must weigh the pleasures to be had by diving against an increased risk of death or injury due to the individual's medical condition. As with any recreational activity, there are no data for diving enabling the calculation of an accurate mathematical probability of injury. Experience and physiological principles only permit a qualitative assessment of relative risk.

For the purposes of this document, **Severe Risk** implies that an individual is believed to be at substantially elevated risk of decompression sickness, pulmonary or otic barotrauma or altered consciousness with subsequent drowning, compared with the general population. The consultants involved in drafting this document would generally discourage a student with such medical problems from diving. **Relative Risk** refers to a moderate increase in risk, which in some instances may be acceptable. To make a decision as to whether diving is contraindicated for this category of medical problems, physicians must base their judgement on an assessment of the individual patient. Some medical problems which may preclude diving are **temporary** in nature or responsive to treatment, allowing the student to dive safely after they have resolved.

Diagnostic studies and specialty consultations should be obtained as indicated to determine the diver's status. A list of references is included to aid in clarifying issues that arise. Physicians and other medical professionals of the Divers Alert Network (DAN) associated with Duke University Health System are available for consultation by phone +1 919 684 2948 during normal business hours. For emergency calls, 24 hours 7 days a week, call +1 919 684 8111 or +1 919 684 4DAN (collect). Related organizations exist in other parts of the world - DAN Europe in Italy +39 039 605 7858, DAN S.E.A.P. in Australia +61 3 9886 9166 and Divers Emergency Service (DES) in Australia +61 8 8212 9242, DAN Japan +81 33590 6501 and DAN Southern Africa +27 11 242 0380. There are also a number of informative websites offering similar advice.

NEUROLOGICAL

Neurological abnormalities affecting a diver's ability to perform exercise should be assessed according to the degree of compromise. Some diving physicians feel that conditions in which there can be a waxing and waning of neurological symptoms and signs, such as migraine or demyelinating disease, contraindicate diving because an exacerbation or attack of the preexisting disease (e.g.: a migraine with aura) may be difficult to distinguish

from neurological decompression sickness. A history of head injury resulting in unconsciousness should be evaluated for risk of seizure.

Relative Risk Conditions

- Complicated Migraine Headaches whose symptoms or severity impair motor or cognitive function, neurologic manifestations
- History of Head Injury with sequelae other than seizure
 Herniated Nucleus Pulposus
- Intracranial Tumor or Aneurysm
- Peripheral Neuropathy
- Multiple Sclerosis
- Trigeminal Neuralgia
- · History of spinal cord or brain injury

Temporary Risk Condition

History of cerebral gas embolism without residual where pulmonary air trapping has been excluded and for which there is a satisfactory explanation and some reason to believe that the probability of recurrence is low.

Severe Risk Conditions

Any abnormalities where there is a significant probability of unconsciousness, hence putting the diver at increased risk of drowning. Divers with spinal cord or brain abnormalities where perfusion is impaired may be at increased risk of decompression sickness.

Some conditions are as follows:

- History of seizures other than childhood febrile seizures
 History of Transient Ischemic Attack (TIA) or Cerebrovascular Accident (CVA)
- History of Serious (Central Nervous System, Cerebral or Inner Ear) Decompression Sickness with residual deficits

CARDIOVASCULAR SYSTEMS

Relative Risk Conditions

The diagnoses listed below potentially render the diver unable to meet the exertional performance requirements likely to be encountered in recreational diving. These conditions may lead the diver to experience cardiac ischemia and its consequences. Formalized stress testing is encouraged if there is any doubt regarding physical performance capability. The suggested minimum criteria for stress testing in such cases is at least 13 METS.* Failure to meet the exercise criteria would be of significant concern. Conditioning and retesting may make later qualification possible. Immersion in water causes a redistribution of blood from the periphery into the central compartment, an effect that is greatest in cold water. The marked increase in cardiac preload during immersion can precipitate pulmonary edema in patients with impaired left ventricular function or significant valvular disease. The effects of immersion can mostly be gauged by an assessment of the diver's performance while swimming on the surface. A large proportion of scuba diving deaths in North America are due to coronary artery disease. Before being approved to scuba dive, individuals older than 40 years are recommended to undergo risk assessment for coronary artery disease. Formal exercise testing may be needed to assess the risk.

* METS is a term used to describe the metabolic cost. The MET at rest is one, two METS is two times the resting level, three METS is three times the resting level, and so on. The resting energy cost (net oxygen requirement) is thus standardized. (Exercise Physiology; Clark, Prentice Hall, 1975.)

Relative Risk Conditions

- History of Coronary Artery Bypass Grafting (CABG)
 Percutaneous Balloon Angioplasty (PCTA) or Coronary Artery Disease (CAD)
- · History of Myocardial Infarction
- Congestive Heart Failure
- Hypertension
- History of dysrythmias requiring medication for suppression
- Valvular Regurgitation

Pacemakers

The pathologic process that necessitated should be addressed regarding the diver's fitness to dive. In those instances where the problem necessitating pacing does not preclude diving, will the diver be able to meet the performance criteria?

* NOTE: Pacemakers must be certified by the manufacturer as able to withstand the pressure changes involved in recreational diving.

Severe Risks

Venous emboli, commonly produced during decompression, may cross major intracardiac right-to-left shunts and enter the cerebral or spinal cord circulations causing neurological decompression illness. Hypertrophic cardiomyopathy and valvular stenosis may lead to the sudden onset of unconsciousness during exercise.

PULMONARY

Any process or lesion that impedes airflow from the lungs places the diver at risk for pulmonary overinflation with alveolar rupture and the possibility of cerebral air embolization. Many interstitial diseases predispose to spontaneous pneumothorax: Asthma (reactive airway disease), Chronic Obstructive Pulmonary Disease (COPD), cystic or cavitating lung diseases may all cause air trapping. The 1996 Undersea and Hyperbaric Medical Society (UHMS) consensus on diving and asthma indicates that for the risk of pulmonary barotrauma and decompression illness to be acceptably low, the asthmatic diver should be asymptomatic and have normal spirometry before and after an exercise test. Inhalation challenge tests (e.g.: using histamine, hypertonic saline or methacholine) are not sufficiently standardized to be interpreted in the context of scuba diving.

A pneumothorax that occurs or reoccurs while diving may be catastrophic. As the diver ascends, air trapped in the cavity expands and could produce a tension pneumothorax.

In addition to the risk of pulmonary barotrauma, respiratory disease due to either structural disorders of the lung or chest wall or neuromuscular disease may impair exercise performance. Structural disorders of the chest or abdominal wall (e.g.: prune belly), or neuromuscular disorders, may impair cough, which could be life threatening if water is aspirated. Respiratory limitation due to disease is compounded by the combined effects of immersion (causing a restrictive deficit) and the increase in gas density, which increases in proportion to the ambient pressure (causing increased airway resistance). Formal exercise testing may be helpful.

Relative Risk Conditions

History of Asthma or Reactive Airway Disease (RAD)*
 History of Exercise Induced Bronchospasm (EIB)*
 History of solid, cystic or cavitating lesion*

- Pneumothorax secondary to:
 - -Thoracic Surgery
 - -Trauma or Pleural Penetration*
 - -Previous Overinflation Injury*

- Obesity
- History of Immersion Pulmonary Edema Restrictive Disease*
- Interstitial lung disease: May increase the risk of pneumothorax
- * Spirometry should be normal before and after exercise

Active Reactive Airway Disease, Active Asthma, Exercise Induced Bronchospasm, Chronic Obstructive Pulmonary Disease or history of same with abnormal PFTs or a positive exercise challenge are concerns for diving.

Severe Risk Conditions

- History of spontaneous pneumothorax. Individuals who
 have experienced spontaneous pneumothorax should avoid
 diving, even after a surgical procedure designed to prevent
 recurrence (such as pleurodesis). Surgical procedures either
 do not correct the underlying lung abnormality (e.g.: pleurodesis, apical pleurectomy) or may not totally correct it (e.g.: resection of blebs or bullae).
- Impaired exercise performance due to respiratory disease.

GASTROINTESTINAL

Temporary Risks

As with other organ systems and disease states, a process which chronically debilitates the diver may impair exercise performance. Additionally, dive activities may take place in areas remote from medical care. The possibility of acute recurrences of disability or lethal symptoms must be considered.

Temporary Risk Conditions

- Peptic Ulcer Disease associated with pyloric obstruction or severe reflux
- Unrepaired hernias of the abdominal wall large enough to contain bowel within the hernia sac could incarcerate.

Relative Risk Conditions

- Inflammatory Bowel Disease
- Functional Bowel Disorders

Severe Risks

Altered anatomical relationships secondary to surgery or malformations that lead to gas trapping may cause serious problems. Gas trapped in a hollow viscous expands as the divers surfaces and can lead to rupture or, in the case of the upper GI tract, emesis. Emesis underwater may lead to drowning.

Severe Risk Conditions

- Gastric outlet obstruction of a degree sufficient to produce recurrent vomiting
- · Chronic or recurrent small bowel obstruction ·

Severe gastroesophageal reflux

- Achalasia
- Paraesophageal Hernia

ORTHOPAEDIC

Relative impairment of mobility, particularly in a boat or ashore with equipment weighing up to 18 kgs/40 pounds must be assessed. Orthopaedic conditions of a degree sufficient to impair exercise performance may increase the risk.

Relative Risk Conditions

- Amputation
- Scoliosis must also assess impact on respiratory function and exercise performance.
- Aseptic Necrosis possible risk of progression due to effects of decompression (evaluate the underlying medical

cause of decompression may accelerate/escalate the progression).

Temporary Risk Conditions

Back pain

HEMATOLOGICAL

Abnormalities resulting in altered rheological properties may theoretically increase the risk of decompression sickness. Bleeding disorders could worsen the effects of otic or sinus barotrauma, and exacerbate the injury associated with inner ear or spinal cord decompression sickness. Spontaneous bleeding into the joints (e.g.: in hemophilia) may be difficult to distinguish from decompression illness.

Relative Risk Conditions

- Sickle Cell Disease
- Polycythemia Vera
- Leukemia
- · Hemophilia/Impaired Coagulation

METABOLIC AND ENDOCRINOLOGICAL

With the exception of diabetes mellitus, states of altered hormonal or metabolic function should be assessed according to their impact on the individual's ability to tolerate the moderate exercise requirement and environmental stress of sport diving. Obesity may predispose the individual to decompression sickness, can impair exercise tolerance and is a risk factor for coronary artery disease.

Relative Risk Conditions

- Hormonal Excess or Deficiency
- Obesity
- Renal Insufficiency

Severe Risk Conditions

The potentially rapid change in level of consciousness associated with hypoglycemia in diabetics on insulin therapy or certain oral hypoglycemic medications can result in drowning. Diving is therefore generally contraindicated, unless associated with a specialized program that addresses these issues.

Pregnancy: The effect of venous emboli formed during decompression on the fetus has not been thoroughly investigated. Diving is therefore not recommended during any stage of pregnancy or for women actively seeking to become pregnant.

BEHAVIORAL HEALTH

Behavioral: The diver's mental capacity and emotional make-up are important to safe diving. The student diver must have sufficient learning abilities to grasp information presented to him by his instructors, be able to safely plan and execute his own dives and react to changes around him in the underwater environment. The student's motivation to learn and his ability to deal with potentially dangerous situations are also crucial to safe scuba diving.

Relative Risk Conditions

- Developmental delay
- · History of drug or alcohol abuse
- · History of previous psychotic episodes
- Use of psychotropic medications

Severe Risk Conditions

 Inappropriate motivation to dive - solely to please spouse, partner or family member, to prove oneself in the face of

personal fears

- · Claustrophobia and agoraphobia
- Active psychosis
- · History of untreated panic disorder
- · Drug or alcohol abuse

OTOLARYNGOLOGICAL

Equalisation of pressure must take place during ascent and descent between ambient water pressure and the external auditory canal, middle ear and paranasal sinuses. Failure of this to occur results at least in pain and in the worst case rupture of the occluded space with disabling and possible lethal consequences.

The inner ear is fluid filled and therefore noncompressible. The flexible interfaces between the middle and inner ear, the round and oval windows are, however, subject to pressure changes. Previously ruptured but healed round or oval window membranes are at increased risk of rupture due to failure to equalise pressure or due to marked overpressurisation during vigorous or explosive Valsalva manoeuvres.

The larynx and pharynx must be free of an obstruction to airflow. The laryngeal and epiglotic structure must function normally to prevent aspiration.

Mandibular and maxillary function must be capable of allowing the patient to hold a scuba mouthpiece. Individuals who have had mid-face fractures may be prone to barotrauma and rupture of the air filled cavities involved.

Relative Risk Conditions

- · Recurrent otitis externa
- Significant obstruction of external auditory canal

History of significant cold injury to pinna

- · Eustachian tube dysfunction
- · Recurrent otitis media or sinusitis
- History of TM perforation
- · History of tympanoplasty
- · History of mastoidectomy
- Significant conductive or sensorineural hearing impairment
- Facial nerve paralysis not associated with barotrauma
 Full prosthedontic devices
- · History of mid-face fracture
- Unhealed oral surgery sites
- History of head and/or neck therapeutic radiation History of temperomandibular joint dysfunction •

History of round window rupture

Severe Risk Conditions

- Monomeric TM
- Open TM perforation
- Tube myringotomy
- · History of stapedectomy
- · History of ossicular chain surgery
- · History of inner ear surgery
- Facial nerve paralysis secondary to barotrauma
 Inner ear disease other than presbycusis
- · Uncorrected upper airway obstruction
- Laryngectomy or status post partial laryngectomy
 Tracheostomy
- Uncorrected laryngocele
- · History of vestibular decompression sickness

BIBLIOGRAPHY/REFERENCE

- Bennett, P. & Elliott, D (eds.)(1993). The Physiology and Medicine of Diving. 4th Ed., W.B. Saunders Company Ltd., London, England.
- Bove, A., & Davis, J. (1990). Diving Medicine. 2nd Edition, W.B. Saunders Company, Philadelphia, PA.
- 3. Davis, J., & Bove, A. (1986). "Medical Examination of Sport Scuba Divers, Medical Seminars, Inc.," San Antonio, TX
- 4. Dembert, M. & Keith, J. (1986). "Evaluating the Potential Pediatric Scuba Diver." AJDC, Vol. 140, November.
- Edmonds, C., Lowry, C., & Pennefether, J. (1992) .3rd ed., Diving and Subaquatic Medicine. Butterworth & Heineman Ltd., Oxford, England.
- Elliott, D. (Ed) (1994). " Medical Assessment of Fitness to Dive." Proceedings of an International Conference at the Edinburgh Conference Centre, Biomedical Seminars, Surry, England.
- "Fitness to Dive," Proceedings of the 34th Underwater & Hyperbaric Medical Society Workshop (1987) UHMS Publication Number 70(WS-FD) Bethesda, MD.

- Neuman, T. & Bove, A. (1994). "Asthma and Diving. Ann. Allergy, Vol. 73, October, O'Conner & Kelsen.
- Shilling, C. & Carlston, D. & Mathias, R. (eds) (1984). The Physician's Guide to Diving Medicine. Plennum Press, New York, NY
- Undersea and Hyperbaric Medical Society (UHMS) www.UHMS.org
- Divers Alert Network (DAN) United States, 6 West Colony Place, Durham, NC www.DiversAlertNetwork.org
- Divers Alert Network Europe, P.O. Box 64026 Roseto, Italy, telephone non-emergency line: weekdays office hours +39-085-893-0333, emergency line 24 hours: +39-039-605-7858
- Divers Alert Network S.E.A.P., P. O. Box 384, Ashburton, Australia, telephone 61-3-9886-9166
- 14. Divers Emergency Service, Australia, www.rah.sa.gov.au/hyperbaric, telephone 61-8-8212-9242
- South Pacific Underwater Medicine Society (SPUMS), P.O. Box 190, Red Hill South, Victoria, Australia, www.spums.org.au
- 16. European Underwater and Baromedical Society, www.eubs.org

ENDORSERS

Paul A. Thombs, M.D., Medical Director Hyperbaric Medical Center St. Luke's Hospital, Denver, CO, USA

Peter Bennett, Ph.D., D.Sc. Professor, Anesthesiology Duke University Medical Center Durham, NC, USA pbennett@dan.duke.edu

Richard E. Moon, M.D., F.A.C.P., F.C.C.P. Departments of Anesthesiology and Pulmonary Medicine Duke University Medical Center Durham, NC, USA

Roy A. Myers, M.D. MIEMS Baltimore, MD, USA

William Clem, M.D., Hyperbaric Consultant Division Presbyterian/St. Luke's Medical Center Denver, CO, USA

John M. Alexander, M.D. Northridge Hospital Los Angeles, CA, USA

Des Gorman, B.Sc., M.B.Ch.B., F.A.C.O.M., F.A.F.O.M., Ph.D.
Professor of Medicine
University of Auckland, Auckland, NZ
d.gorman@auckland.ac.nz

Alf O. Brubakk, M.D., Ph.D.

Norwegian University of Science and Technology
Trondheim, Norway
alfb@medisin.ntnu.no

Alessandro Marroni, M.D. Director, DAN Europe Roseto, Italy Hugh Greer, M.D. Santa Barbara, CA, USA hdqblqfpl@aol.com Christopher J. Acott, M.B.B.S., Dip. D.H.M., F.A.N.Z.C.A. Physician in Charge, Diving Medicine Royal Adelaide Hospital Adelaide, SA 5000, Australia

Chris Edge, M.A., Ph.D., M.B.B.S., A.F.O.M. Nuffield Department of Anaesthetics Radcliffe Infirmary Oxford, United Kingdom cjedge@diver.demon.co.uk

Richard Vann, Ph.D. Duke University Medical Center Durham, NC, USA

Keith Van Meter, M.D., F.A.C.E.P. Assistant Clinical Professor of Surgery Tulane University School of Medicine New Orleans, LA, USA

Robert W. Goldmann, M.D. St. Luke's Hospital Milwaukee, WI, USA

Paul G. Linaweaver, M.D., F.A.C.P. Santa Barbara Medical Clinic Undersea Medical Specialist Santa Barbara, CA, USA

James Vorosmarti, M.D. 6 Orchard Way South Rockville, MD, USA

Tom S. Neuman, M.D., F.A.C.P., F.A.C.P.M. Associate Director, Emergency Medical Services Professor of Medicine and Surgery University of California at San Diego San Diego, CA, USA

Yoshihiro Mano, M.D. Professor Tokyo Medical and Dental University Tokyo, Japan y.mano.ns@tmd.ac.jp Simon Mitchell, MB.ChB., DipDHM, Ph.D. Wesley Centre for Hyperbaric Medicine Medical Director Sandford Jackson Bldg., 30 Chasely Street Auchenflower, QLD 4066 Australia smitchell@wesley.com.au

Jan Risberg, M.D., Ph.D. NUI, Norway

Karen B.Van Hoesen, M.D. Associate Clinical Professor UCSD Diving Medicine Center University of California at San Diego San Diego, CA, USA

Edmond Kay, M.D., F.A.A.F.P. Dive Physician & Asst. Clinical Prof. of Family Medicine University of Washington Seattle, WA, USA ekay@u.washington.edu

Christopher W. Dueker, TWS, M.D. Atherton, CA, USA chrisduek@aol.com

Charles E. Lehner, Ph.D.
Department of Surgical Sciences
University of Wisconsin
Madison, WI, USA
celehner@facstaff.wisc.edu

Undersea & Hyperbaric Medical Society 10531 Metropolitan Avenue Kensington, MD 20895, USA

Diver's Alert Network (DAN) 6 West Colony Place Durham, NC 27705